

- Gass, C. L. (1978), 'Territory regulation, tenure, and migration in Rufous Hummingbirds', *Can. J. Zool.* 57: 914-923.
- des Granges, J. L. (1979), 'Organization of a tropical nectar-feeding bird guild in a variable environment', *Living Bird* 17: 199-236.
- Hainsworth, F. R. and L. L. Wolf (1979), 'Feeding: an ecological approach', *Adv. Stud. Behav.* 9: 53-96.
- Immelmann, K. (1961), 'A contribution to the biology and ethology of Australian honeyeaters', *J. für Ornithologie* 102: 164-207.
- Kikkawa, J., I. Hore-Lacy and J. le Gay Brereton (1965), 'A preliminary report on the birds of the New England National Park', *Emu* 65: 139-143.
- Paton, D. C. (1979), 'The behaviour and feeding ecology of the New Holland Honeyeater, *Phylidonyris novaehollandiae*, in Victoria', Ph.D. thesis, Monash University, Melbourne.
- Pyke, G. (1980), 'The foraging behaviour of Australian honeyeaters: a review and some comparisons with hummingbirds', *Aust. J. Ecol.* 5: 343-369.
- Rooke, I. (1979), 'The social behaviour of the honeyeater *Phylidonyris novaehollandiae*', Ph.D. thesis, University of Western Australia. RAOU Microfiche Series No. 3, Melbourne.
- Stiles, F. G. and L. L. Wolf (1970), 'Hummingbird territoriality in a flowering tropical tree', *Auk* 87: 467-491.
- Thomas, D. G. (1980), 'Foraging of honeyeaters in an area of Tasmanian sclerophyll forest', *Emu* 80: 55-58.
- Tullis, K. J. and R. D. Wooller (1981), 'Interspecific feeding hierarchies in captive honeyeaters', *Bird Behaviour* 3: 93-95.

Corella, 1985, 9: 25-29

## A SURVEY OF PELAGIC BIRDS IN THE WESTERN CORAL SEA AND GREAT BARRIER REEF

TONY STOKES\* and CHRIS CORBEN†

\* Government Conservator, Australian National Parks and Wildlife Service, Christmas Island, Indian Ocean 6798.

† 26, Warmington Street, Paddington, Queensland, 4064.

Received 5 March, 1984

Little is known of the distribution of seabirds in the Western Coral Sea, i.e. the area west of 155°E longitude and north of 25°S latitude nominated by Pickard *et al.* (1977). Most of the area is within the Coral Sea Islands Territory of Australia which extends west to 157°10' longitude, north to 12° and south to 24° latitude.

Although Serventy (1959) summarised what is known of the birds of Willis Island, where there is a manned lighthouse, many of the small cays and islands have only been visited once by ornithologists, in 1961/1962 (Hindwood *et al.* 1963). In the winters of the same years Norris (1967) made the most comprehensive observations yet available on pelagic birds in the area.

The cays and islands in Figure 1 have automatic weather stations or lighthouse installations and have been surveyed at least once and usually twice per year since 1979 by the Australian National Parks and Wildlife Service (ANPWS) which is developing a nature conservation programme for the Coral Sea Islands Territory. The results discussed in this paper were obtained during the course of one survey.

### THE SURVEY

We surveyed a section of the Coral Sea and Great Barrier Reef between 15 to 26 May 1981 (Figure 1). A total of 50 hours 45 minutes was spent on watch from the M.V. *Cape Pillar*.

No weather details were recorded, but although occasional squalls passed over, the watch was usually conducted in rain-free conditions on smooth to moderate seas.

Observations were virtually continuous during daylight steaming. Sightings were made from the Flying Bridge about 15 m above the sea surface. 8 x 40 binoculars were used by Stokes and 10 x 50 by Corben. Identifications were principally by Corben. Field notes on unidentified species were taken. These notes were later checked with museum skins, photographs and text descriptions and the record amended where necessary. Though every effort was made not to recount birds that disappeared from view and reappeared elsewhere it is probable that some were recounted.

The seabird records are summarised in Table 1 and Figure 1. The sightings are based on the ship's position by 'Dead Reckoning'. No allowance is made for the distance of the bird from the ship.

Figure 1 is plotted onto a 30' grid of latitude and longitude. The relevant grid squares have been identified by letters which are used in Table 1. The duration of observations in each grid is given in Table 1.

The scientific names of species mentioned in the text are in Table 1. The names of cays are those used by Hindwood *et al.* (1963).

## DISCUSSION

Many of the observations of large numbers of a species in a grid square relate to congregations on nearby cays or islands some of which are identified in Figure 1.

Two exceptions are the flocks of birds seen in squares j, k and l near the edge of the Great Barrier Reef, and the large numbers of Hutton's or Fluttering Shearwaters in the vicinity of Gannet Cay.

The more interesting records are discussed below.

The Herald Petrel has been recorded on a few occasions in the western Coral Sea and eastern Australian waters in January, February and May (Izzard and Watson 1980). Corben also saw one fly over Turtle Islet (grid 'z') during this trip in




Figure 1. Map showing locations of reefs and cays referred to in text.

daylight on 24 May 1981 and the bird must be considered a regular but rare species in the area.

Sightings of Tahiti Petrels were concentrated in grid squares outside the Great Barrier Reef. Norris (1967) and Hindwood *et al.* (1963) also recorded the species from western Coral Sea waters in July and October to November. Holmes (1981a) identifies the Hindwood *et al.* (1963, p.8) unidentified records as belonging to this species and lists more recent records from central eastern Australian waters. He interprets oceanographic data to imply that the species moves south with warm Coral Sea waters in January and February.

Certainly the Tahiti Petrel appears to be relatively common over western Coral Sea waters



from May to November and many of the birds would presumably originate from breeding colonies on Chesterfield Reef islands in the eastern Coral Sea. Further observations are necessary to distinguish the relative numbers and distribution in the area of the two forms identified by Holmes (1981a). We did not distinguish forms.

Little is known of the dispersal of Gould's Petrel from its breeding grounds (Serventy *et al.* 1971, Pizzey 1980). The nominate subspecies breeds only on Cabbage Tree Island, New South Wales, and there are regular sightings and beach-washed specimens from central to eastern Australia between November to March, its breeding season (Ingram 1975, Gosper 1981, Holmes 1981b). Gibson and Sefton (1957) thought that the species "... shows a predilection for warmer seas in the winter months" which Cox (1976) supported. However Pizzey (1980) admits that its movements are little known and there are few records outside the breeding season.

A different subspecies of Gould's Petrel breeds on New Caledonia from where it appears to forage in the Tasman Sea and migrates to the south-west Pacific Ocean (Imber and Jenkins 1981). It is possible that breeding birds also forage in the Coral Sea.

A fresh Sooty Shearwater carcass was found on Gannet Cay on 19 May. It and the bird at sea if correctly identified, would have been migrating north (Phillips 1963, Serventy *et al.* 1971).

The movements of Wedge-tailed Shearwaters from their breeding islands off eastern Australia are uncertain but there are indications that the population from the coastal islands of New South Wales regularly migrate north, perhaps entirely into the northern hemisphere (Purchase 1974, Rogers 1975, Marchant 1977, Milledge 1977). Norris (1967) suggests that at least some birds from the Coral Sea island colonies do likewise as he recorded few over the Coral Sea in July and September. Studies are required to determine the composition and status of birds in the area from May to September.

Our sightings of Hutton's or of Fluttering Shearwaters were concentrated near Swains Reef. Despite extensive comparative research

by Corben, since the voyage, into the plumages of both species it has not been possible to separate with certainty the specimens reported in this paper. Corben *et al.* (1974) first recorded Hutton's Shearwaters from Queensland waters and thought that it was a regular but rare visitor to the south-east coast and accidental in the north. Pizzey (1980) agrees and adds that both species often flock together.

Wilson's Storm-Petrels were seen principally over the Great Barrier Reef supporting the Serventy *et al.* (1971) statement that they are essentially birds of continental shelf waters. The other Storm-Petrels were more common in the northern grid squares over waters outside the Reef. A White-faced Storm-Petrel was seen near Gannet Cay; this is notable for a bird that 'seems to be markedly pelagic in habit' (Serventy *et al.* 1971). The White-bellied Storm-Petrel records are further evidence that it is common in the area (Hindwood *et al.* 1963, Norris 1967, Serventy *et al.* 1971). The Black-bellied Storm Petrel record is somewhat north of the 30°-60°S latitude range which it is said to frequent while wintering off Australia (Serventy *et al.* 1971) and appears to be the northern most record from waters off the Queensland coast (Roberts 1973).

Tree Martins are frequently windswept over the ocean off eastern Australia, apparently while on migration. Apart from the two records in Table 1, one was blown across the unvegetated sandy Cay A, Flinders Reef, on 25 May 1981 and a carcass was found by ANPWS staff on 29 April 1980 on the unvegetated North Reef Cay of Frederick Reef. Serventy (1959) and Warham (1961) also report specimens from the virtually treeless Raine and Willis Islands, respectively.

Many of the remaining species seen during the survey are common breeding birds in the area but little is known of their foraging range or non-breeding dispersal from the islands. However, complex currents operate in the area with a 'flushing' effect on parts of the Great Barrier Reef (Pickard *et al.* 1977) and, with upwellings of cold water, probably affect seabird distribution.

Recently proclaimed nature reserves in the Herald, Coringa and Magdelaine Islands and Lihou Reef will assist the conservation of the birds.

### ACKNOWLEDGEMENTS

We are grateful to the Australian National Parks and Wildlife Service for proposing and funding the survey, and to Captain F. Hanson and the Department of Transport for facilitating it.

Captain M. Carrell and the crew of the M.V. *Cape Pillar* were most hospitable and provided every assistance.

Dr K. Hulsman suggested Mr Corben's participation and provided material for collecting carcasses.

Comments on drafts of the text were kindly provided by John Henry, Keiran McNamara and Vance Russell.

### REFERENCES

- Corben, C. J., G. J. Roberts and G. J. Ingram (1974), 'Sightings of Huttons Shearwater in Queensland', *Sunbird* 5: 55-56.
- Cox, J. B. (1976), 'A review of Procellariiformes occurring in South Australian waters', *South Aust. Orn.* 27: 28-82.
- Gibson, J. D. and A. R. Sefton (1957), 'The Gould Petrel — Australian Records', *Emu* 57: 49-52.
- Gosper, D. G. (1981), 'A survey of beach-washed seabirds near Ballina, New South Wales', *Corella* 5: 110-113.
- Hindwood, K. A., K. Keith and D. L. Serventy (1963), 'Birds of the South-West Coral Sea', CSIRO Div. Wildl. Res. Tech. Paper No. 3.
- Holmes, G. (1981a), 'Occurrence and identification of the Tahiti Petrel in central eastern Australian (sic)', *Aust. Birds* 16: 24.
- Holmes, G. (1981b), 'Unequal sex ratios among seabirds found beachwashed', *Emu* 81: 44-47.
- Imber, M. J. and J. A. F. Jenkins (1981), 'The New Caledonian Petrel', *Notornis* 28: 149-160.
- Ingram, G. J. (1975), 'Records of the Gould Petrel and the Black-winged Petrel from south-eastern Queensland and north-eastern New South Wales', *Sunbird* 6: 92-95.
- Izzard, J. and W. D. Watson (1980), 'A sight record of the Herald Petrel off Northern New South Wales', *Aust. Birds* 15: 5-6.
- Marchant, S. (1977), 'A Seawatch on the southern coast of New South Wales', *Emu* 77: 9-18.
- Milledge, D. R. (1977), 'One year's observations of seabirds in Continental Shelf waters off Sydney, N.S.W.', *Corella* 1: 1-12.
- Norris, A. Y. (1967), 'Seabird observations from the south-west Pacific in the southern winter', *Emu* 67: 33-54.
- Phillips, J. H. (1963), 'The Pelagic Distribution of the Sooty Shearwater *Procellaria grisea*', *Ibis* 105: 340-353.
- Pickard, G. L., J. R. Donguy, C. Henin and F. Rougerie (1977), 'A Review of the Physical Oceanography of the Great Barrier Reef and Western Coral Sea', *Aust. Inst. Mar. Sci. Mon.* 2.
- Pizzey, G. (1980), *A Field Guide to the Birds of Australia*. Collins: Sydney.
- Purchase, D. (1974), 'First overseas recoveries of Wedge-tailed Shearwaters banded in Australia', *Aust. Bird Bander* 12: 34-35.
- Roberts, G. (1973), 'Specimen records of the Black-bellied Storm-petrel from South-east Queensland', *Sunbird* 4: 52-53.
- Rogers, A. E. F. (1975), 'N.S.W. Bird Report for 1974', *Aust. Birds* 9: 77-97.
- Serventy, D., V. Serventy and J. Warham (1971), *The Handbook of Australian Seabirds*. A. H. & A. W. Reed: Sydney.
- Serventy, V. (1959), 'Birds of Willis Island', *Emu* 59: 167-176.
- Warham, J. (1961), 'The birds of Raine Island, Pandora Cay and Murray Island Sandbank, north Queensland', *Emu* 61: 77-93.