


**Rufous-banded Honeyeater *Conopophila albogularis* Species No.: 600 Band size: 02**



**Morphometrics:**

	Adult Male	Adult Female
Wing:	62 – 69 mm	59 – 66 mm
Wing Span:	> 216 mm	< 216 mm
Tail:	41 – 50 mm	38 – 48 mm
Weight:	9.2 – 14.5 g	9.8 – 13.2 g

**Ageing:**

	Adult (1+)	Juvenile
Bill:	greyish-black to black with paler bluish-grey base to lower mandible;	grey-black with pinkish, salmon, flesh-coloured or orange base to lower mandible;
Gape:	pale grey, or bluish grey;	yellow, salmon, pinkish or orange;
Iris:	brown, brownish-grey, red, light brown or olive-brown;	grey or olive-grey;
Forehead:	grey with brownish tinge to front;	brownish-grey, browner posteriorly;
Crown & nape:	grey with dark brown streaking	grey-brown or brown;
Lores:	grey or dark grey;	brownish-grey;
Malar area:	grey;	pale grey;
Chin & throat:	white;	off-white grading to light brown on sides;
Upper breast:	cinnamon brown forming a distinct band;	off-white or pale buff

Juveniles moult to adult-like plumage within a few months of fledging;  
It has not been documented as to whether immatures retain juvenile plumage such as alula, remiges and rectrices, but, like other honeyeaters, it is likely that some or all are retained until a complete moult commencing at about 1 year old;  
Because of lack of information adults should only be aged (1+).

**Sexing :**

Sexes do not differ in plumage, but males average larger than females;  
Incubation by female only.