
Yellow-tinted Honeyeater Lichenostomus flavescens Species No.: 624 Band size: 02



Morphometrics:

Adult Male	Adult Female
> 32.7 mm	< 30.6 mm
67 – 81 mm	65 – 73 mm
54 – 66 mm	53 – 64 mm
11 – 16 g	9.4 – 15.5 g
	67 – 81 mm 54 – 66 mm

Ageing:

Adult (1+) Juvenile

Bill: black; yellow or brown with black tip, but becomes

all black within the first year;

Gape: grey or black; yellow;

Iris: dark brown, red-brown, olive-brown grey or dark brown;

or grey-brown;

Forehead & yellow-olive; pale yellowish olive;

forecrown:

Hindcrown & light grey-brown or olive-grey with light grey-brown with faint olive tinge;

nape: yellowish-olive suffusion;

Breast: pale yellow with diffuse light grey- off-white with pale yellow suffusion;

brown streaking;

Juveniles undergo a partial moult to immature plumage that basically resembles adult plumage soon after fledging, but immatures may retain some or all juvenile primaries;

Presence of synchronous growth or fault bars in the tail indicate retained juvenile feathers – thus immature aged (1);

Adult plumage is acquired in the first year, so all adult plumaged birds through the breeding season (late Winter to early Summer) are aged (2+), otherwise (1+);

Sexing:

No sexual dimorphism in plumage, but males larger than females (see THL and wing length above); Not known whether both sexes incubate.