
Emerald Dove Chalcophaps indica Species No.: 33 Band size: 07



Morphometrics:

C.i. chrysochlora	C.i. longirostris
C.I. CIII y SOCIIIOI a	C.I. IOIIGII OSLI IS

	Adult Male	Adult Female	Adult Male	Adult Female
Wing:	146 – 161 mm	143 – 152 mm	153 – 165 mm	151 – 160 mm
Tail:	78 – 90 mm	75 – 88 mm	82 – 94 mm	80 – 90 mm
Weight:	120 – 172 g	124 – 155 g	140 – 192 g	150 – 167 g

Ageing:

Adult (1+) Juvenile

Bill: orange-yellow with purplish-red black-brown with cream tip;

base;

Iris: dark brown; black-brown; Orbital skin: pink-red; dark brown;

Legs: dull red to dark purplish-red; dark grey with red tinge;

Shoulder patch: see below; absent;

Median & greater iridescent dark-green; dark brown with rufous tips;

secondary coverts:

Complete post-juvenile moult to adult plumage has finished at about 3 months, thus adults should be aged (1+)

Sexing:

Adults can be sexed on plumage as follows:

Adult Male Adult Female

Crown & nape dirty pink with reddish-brown tinge; dark brown with broad dark red-brown

fringes;

Chin: pinkish buff to off-white; light rufous-brown; Rump & upper black-brown with broad black red-brown to dark red-brown

tail coverts: fringes to feathers:

Breast & belly: dirty pink with reddish-brown tinge brown to light rufous-brown

grading to pink brown on belly;

Shoulder patch: lesser secondary coverts mostly

coverts: grey with brown bases, but some

larger secondary coverts have broad white fringes forming

prominent white shoulder patch;

brown to light rufous-brown grading to light grey-brown on belly;

mostly light brown with larger secondary coverts having narrow light-grey to pale -grey tips forming an inconspicuous

grey shoulder patch;

Role of sexes in incubation is not known