Double-eyed Fig-Parrot Cyclopsitta diophthalma Species No.: 261 Band size: 24 Adult Male

Female

ssp. C.d. marshalli

Adult

Female

ssp. C.d. macleayana

ssp. C.d. coxeni

Male

Adult

Female

Morphometrics:

Three subspecies in Australia and morphometrics are as follows:

Juvenile

C.d. marshalli (Cape York, Qld);	C.d. macleayana (South of
	Cooktown to Townsville Old):

Adult

Male

			COURTOWN TO TOWNSVINE, QIU),	
	Adult Male	Adult Female	Adult Male	Adult Female
Wing:	81 – 89 mm	79 – 89 mm	86 – 94 mm	43 – 47 mm
Tail:	40 – 46 mm	37 – 43 mm	43 – 47 mm	41 – 46 mm
Bill:	12.9 – 14.3 mm	12.9 – 14.2 mm	12.7 – 14.3 mm	13.1 – 14.3 m
Tarsus:	11.5 – 12.3 mm	11.8 – 12.7 mm	12.1 – 13.2 mm	11.5 – 13.4 mm
Weiaht:	25 – 42 a	35 – 41 a	few re	ecords

C.d. coxeni (s-e Qld & n-e NSW)

	Adult Male	Adult Female	
Wing:	96 – 101 mm	95 – 97 mm	
Tail:	44 – 48 mm	46 – 47 mm	
Bill:	15.5 – 16.2 mm	15.3 -15.7 mm	
Tarsus:	13.3 – 14.3 mm	14.1 – 15.0 mm	
Weight:	no records		

Ageing:

Juveniles and immatures of *C.d. marshalli* and *C.d. macleayana* are duller than, but difficult to distinguish from adult females. Age all female plumaged birds of these subspecies as (1+);

Adult male plumage is attained at approximately 14 months. Thus age all adult males of these two subspecies as (2+).

Female adults of *C.d. coxeni* are said to be slightly duller than males, but can not be reliably sexed on plumage. Therefore age all *coxeni* as (1+).

Sexing:

See illustrations above.